



BIOCHEMICAL MODIFICATIONS IN FRUIT MATURATION: INVESTIGATING CHEMICAL PATHWAYS AND RIPENING MECHANISMS

Sharvesh S¹, Sharmitha K², Dharanidharan S³ and Sharmila B^{4*}

¹Research scholar, Department of Horticulture, Faculty of Agriculture, Annamalai University, Chidambaram

²Assistant Professor, Prist univeristy, Thanjavur

³Assistant Professor, Aravindhar Agricultural Institute of Technology,

⁴Assistant Professor, AVS College of Technology, Salem

*Corresponding Author Mail ID: samsherin98@gmail.com

Introduction

Fruits undergo complex chemical transformations during maturation and ripening, processes that are critical to determining their quality, texture, taste and overall appeal. These changes, regulated by both internal metabolic pathways and external factors, ensure that a fruit progresses from being inedible and hard to a soft, sweet and flavorful state. Understanding these processes is crucial not only for scientific study but also for practical applications like improving post-harvest handling, storage and marketing strategies.

Maturation vs Ripening

Before delving into the specific chemical changes, it is important to differentiate between maturation and ripening. Maturation is the phase during which the fruit is still on the plant, undergoing physiological development to reach the stage at which it can continue ripening after harvest. Maturation sets the stage for ripening, which is the final phase where significant biochemical changes occur, turning the fruit into its edible form.

Maturity: This refers to the developmental stage where a fruit can attain the desired quality after harvest. Maturity is critical because fruits harvested too early or too late may not ripen properly, reducing their appeal and shelf life.

Ripening: This phase starts after the fruit has matured and is characterized by significant biochemical and physiological changes. These changes involve the transformation of complex

compounds into simpler ones, colour alterations, softening of tissues and the development of characteristic flavours and aromas.

Chemical Changes During Ripening

Several biochemical reactions and pathways govern the ripening process. These transformations affect various aspects of the fruit, including its texture, color, sugar content, acidity and aroma.

Hydrolysis of Polysaccharides to Sugars

One of the most critical changes in fruit ripening is the breakdown of polysaccharides (like starch) into simpler sugars (like glucose, fructose and sucrose). This conversion is what makes the fruit taste sweet and desirable. In starchy fruits like bananas and apples, this breakdown is particularly pronounced and occurs rapidly as ripening progresses.

During maturation, fruits typically store energy in the form of polysaccharides. As the ripening process begins, enzymes such as amylase and glucanases catalyze the hydrolysis of starch into sugars. The increase in sugar content is associated with the development of a more pleasant taste and a softer texture, making the fruit more palatable.

Reduction in Acidity

The acidity in fruits typically decreases as ripening progresses, which contributes to the sweet taste of ripe fruits. Organic acids, such as malic acid, citric acid and oxalic acid, present in fruits serve as a reserve of metabolic energy. During ripening, these acids are either converted

into sugars or metabolized in the respiration process, leading to a reduction in the overall acidity of the fruit.

This decline in acidity helps achieve the sugar-acid balance that is crucial for the flavour profile of the fruit. For example, fruits like tomatoes and citrus, which initially have high levels of organic acids, undergo significant flavour improvements as their acidity diminishes during ripening.

Chlorophyll Degradation and Colour Changes

One of the most noticeable changes during fruit ripening is the change in colour. Unripe fruits are typically green due to the presence of chlorophyll, a pigment responsible for photosynthesis. As the fruit ripens, chlorophyll breaks down, a process termed degreening and other pigments like carotenoids (yellow to orange) and anthocyanins (red to purple) become more visible.

This change in colour is significant not only as a visual cue for ripeness but also for the fruit's appeal. For example, the deepening of red in tomatoes or strawberries and the yellowing of bananas signal that the fruit is ready to eat. The breakdown of chlorophyll is facilitated by chlorophyllase, which degrades the magnesium-chlorophyll complex, allowing other pigments to dominate.

Softening Due to Pectin Breakdown

A critical factor in the texture of fruit is its firmness, which decreases as the fruit ripens. This softening is largely due to the breakdown of pectic substances in the cell walls. Pectins are complex carbohydrates that give structure and rigidity to plant tissues. As fruit ripens, pectinase enzymes solubilize and degrade these substances, resulting in a softer texture.

This softening process is essential for making the fruit easier to consume but must be carefully controlled during storage and transportation to prevent excessive softening, which can lead to spoilage.

Ethylene Production and Respiration Changes

Ethylene, a gaseous plant hormone, plays a central role in regulating fruit ripening,

particularly in climacteric fruits such as bananas, tomatoes and apples. Climacteric fruits exhibit a sharp increase in respiration (known as the respiratory climacteric) and a simultaneous rise in ethylene production during ripening.

Ethylene acts as a signaling molecule, triggering the activation of various enzymes involved in ripening, such as those responsible for colour change, softening and flavour development. The production of ethylene is autocatalytic, meaning that a small amount of ethylene can stimulate the production of more ethylene, accelerating the ripening process.

Non-climacteric fruits, such as strawberries and grapes, do not exhibit this dramatic increase in respiration or ethylene production. Instead, their ripening is usually a more gradual process and is typically completed while still attached to the plant.

Volatile Compound Production and Flavour Development

The characteristic aroma of ripe fruit is due to the production of volatile organic compounds. These compounds are synthesized during the later stages of ripening and include esters, aldehydes, alcohols and ketones. For example, the distinctive aroma of ripe bananas is primarily due to the production of esters like isoamyl acetate and butyl acetate.

Volatile compounds are crucial for the perception of fruit flavour. They contribute to the complex interplay of sweetness, acidity and aroma that defines the fruit's sensory appeal. The exact composition and concentration of these volatiles vary significantly between different types of fruits and are often responsible for the unique flavour profile of each fruit.

Phenolic Compound Change

Phenolic compounds, including tannins, are important for the astringency and bitterness of many unripe fruits. As the fruit ripens, the astringency decreases due to the polymerization of tannins into insoluble forms, making the fruit more palatable.

Additionally, phenolic compounds are responsible for the oxidative browning reaction that occurs when fruits like apples or bananas are

cut. This browning is due to the action of polyphenol oxidase (PPO), which catalyzes the oxidation of phenolics to form brown pigments.

Role of Ethylene in Fruit Ripening

Ethylene, often referred to as the "ripening hormone," is crucial in regulating the entire process of ripening, especially in climacteric fruits. It not only influences the aforementioned changes but also acts as a signaling molecule for gene expression related to ripening. Ethylene regulates:

Softening: By activating enzymes that degrade pectin and other cell wall components.

Colour change: By triggering the degradation of chlorophyll and the synthesis of carotenoids and anthocyanins.

Flavour and aroma: By promoting the production of volatile compounds.

Ethylene can be applied exogenously to control ripening in a post-harvest setting, which is a common practice for fruits like bananas and tomatoes. However, excessive ethylene exposure can lead to over-ripening and spoilage.

Conclusion

The chemical changes that occur during the maturation and ripening of fruits are intricate and involve numerous metabolic pathways. These changes convert unripe fruits into soft, sweet and flavourful products that appeal to consumers. Key processes include the hydrolysis of polysaccharides, reduction in acidity, chlorophyll degradation, pectin breakdown, ethylene production and the synthesis of volatile compounds.

These transformations not only improve the taste, texture and appearance of fruits but also impact their nutritional value and shelf life. Understanding these processes is essential for improving post-harvest handling and ensuring that fruits reach consumers in optimal condition.