



ECO-FRIENDLY WEED MANAGEMENT: A SUSTAINABLE PATH FOR CROP PRODUCTION

Sangothari A¹, Radhamanai S^{2*}, Nagarajan K³ and Bharaani Sri C⁴

¹Senior Research Fellow, Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore-03

²Professor, Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore-03

³Research Associate, Central Institute of Cotton Research, Regional Station, Coimbatore-03

⁴Business Executive, ABIS-TBI, Tamil Nadu Agricultural University, Coimbatore-03

*Corresponding Author Mail ID: radhamani.s@tnau.ac.in

Introduction

Weeds are one of the major hidden enemies of agriculture. They compete with crops for nutrients, water, light and space, ultimately reducing yield and quality. In many cropping systems, yield losses caused by weeds are higher than those caused by insects or diseases. Farmers often spend a large share of cultivation costs on weed control, mainly due to labour shortage and increasing dependence on herbicides.

While chemicals provide quick control, their repeated and indiscriminate use has created serious problems such as herbicide resistance, weed shifts, soil and water pollution and harm to beneficial organisms. These challenges highlight the need for eco-friendly weed management, which focuses on long-term sustainability rather than short-term control.

Why eco-friendly weed management is essential

Eco-friendly weed management aims not at complete weed eradication but at maintaining weed populations at manageable levels. A small weed presence is acceptable if it does not reduce crop yield or quality. This approach helps conserve biodiversity, reduces input costs and protects natural resources. It also aligns with climate-smart agriculture by lowering carbon emissions and energy use associated with intensive chemical weed control.

Approaches in sustainable weed management

1. Cultural methods of weed management

Cultural practices are preventive in nature and play a major role in suppressing weeds naturally.

i. Proper crop stand and plant population

Maintaining optimum plant density through correct seed rate, spacing, sowing time and variety selection helps crops establish quickly and shade weeds. Crops with rapid early growth and dense canopy formation suppress weed emergence effectively.

ii. Green manuring

Green manure crops improve soil fertility and reduce weed infestation. Their rapid growth and biomass production smother weeds, while their incorporation into soil improves nutrient availability and disrupts weed seed survival.

iii. Intercropping

Growing two or more crops together increases crop diversity and resource utilization. Intercropping limits weed growth by reducing open spaces and creating competition for light, moisture and nutrients.

iv. Selection of competitive crop varieties

Crop varieties with early vigor, higher tillering capacity and better canopy structure are more effective in suppressing weeds. Transplanted crops often have a competitive advantage over direct-seeded crops during early growth stages.

v. Crop establishment techniques

Different establishment methods influence weed emergence. Practices such as raised bed planting, zero tillage and furrow-irrigated systems reduce surface moisture availability for weed germination and disturb weed seed banks less frequently.

vi. Planting geometry and row orientation

Closer row spacing and uniform planting reduce sunlight reaching the soil surface, thereby suppressing weed growth. Improved canopy coverage enhances crop dominance over weeds.

vii. Seed rate management

Using appropriate seed rates ensures uniform crop stand and faster ground coverage. Higher plant density often reduces weed biomass through shading and competition.

viii. Water management

Proper irrigation scheduling can discourage weed growth. Some weeds fail to germinate under flooded or saturated conditions, while controlled irrigation reduces favorable conditions for weed establishment.

ix. Cover crops and smother crops

Cover crops protect soil from erosion and suppress weeds by shading and competing for resources. Their residues further act as mulch, reducing weed emergence in subsequent crops.

x. Crop rotation

Rotating crops with different growth habits interrupts weed life cycles and prevents dominance of specific weed species. Crop rotation also reduces weed seed buildup in soil.

xi. Nutrient management

Fertilizer placement and timing significantly affect weed-crop competition. Targeted nutrient application favors crop uptake and limits nutrient availability to weeds.

xii. Use of organic manures

Organic manures improve soil structure, microbial activity and nutrient availability. Certain organic amendments also reduce weed seed viability through biological and chemical interactions in soil.

2. Physical methods of weed management

Physical methods remove weeds using manual or mechanical force and are environmentally safe.

i. Manual and mechanical weeding

Hand weeding, hoeing and mechanical weeders physically uproot weeds, particularly during early growth stages. These methods are effective in reducing weed pressure without chemical residues.

ii. Off-season and summer ploughing

Deep ploughing during summer exposes weed seeds and underground propagules to heat and desiccation, reducing their survival and infestation in the following season.

iii. Soil solarization

Covering moist soil with transparent sheets during hot months raises soil temperature, destroying weed seeds and young seedlings. This method also improves soil health.

iv. Use of weeders and implements

Improved tools such as power weeders, cono weeders and mini tillers reduce labour requirement and drudgery, especially in wide-spaced crops and rice systems.

v. Hand digging and uprooting

Deep-rooted and perennial weeds are effectively controlled by digging and removing underground parts to prevent regrowth.

vi. Burning

Controlled burning destroys weed seeds present on the soil surface but must be used cautiously due to loss of organic matter and environmental concerns.

vii. Mulching

Mulches suppress weeds by blocking sunlight, conserving moisture, regulating soil temperature and improving soil properties. Both organic and biodegradable mulches are effective.

3. Biological weed management

Biological weed management uses living organisms to suppress weeds naturally.

i. Biological control agents

Insects, pathogens and microorganisms specifically target certain weeds, reducing their growth and spread without affecting crops.

ii. Allelopathic plants

Some plants release natural chemicals that inhibit weed germination and growth. These plants can be used as cover crops, mulches, or in rotations.

Table 1. Plants exhibiting allelopathic effects for weed suppression

Allelopathic plant species	Weeds affected / suppressed
<i>Acalypha indica</i>	Broad-leaved weeds
<i>Trianthema portulacastrum</i>	Annual weed species
<i>Eichhornia crassipes</i>	Aquatic and semi-aquatic weeds
<i>Parthenium hysterophorus</i>	Several associated weed species
<i>Cyperus rotundus</i>	Grasses and sedges
<i>Argemone mexicana</i>	Broad-leaved weeds
<i>Lantana camara</i>	Annual and perennial weeds
<i>Eupatorium adenophorum</i>	Competing terrestrial weeds
<i>Imperata cylindrica</i>	Broad-leaved and grassy weeds
<i>Mikania micrantha</i>	Fast-growing annual weeds
<i>Echinochloa colonum</i>	Upland and lowland weeds
<i>Phalaris minor</i>	Grass weeds in cereal systems
<i>Portulaca oleracea</i>	Broad-leaved weeds

iii. Bio-fertilizers in weed suppression

Bio-fertilizers enhance crop vigor and soil microbial activity, indirectly improving crop competitiveness against weeds.

iv. Bio-herbicides

Bio-herbicides are formulations of naturally occurring pathogens that selectively control specific weed species in a sustainable manner.

Table 2. Bio-herbicides and microorganisms used in weed control

Microorganism/pathogen	Target weed species	Crop / ecosystem	Commercial product
<i>Colletotrichum gloeosporioides</i>	<i>Aeschynomene virginica</i>	Rice, soybean	Collego
<i>Phytophthora palmivora</i>	<i>Morrenia odorata</i>	Citrus orchards	DeVine
<i>Alternaria cassiae</i>	<i>Cassia obtusifolia</i>	Soybean	CASST
<i>Alternaria destruens</i>	Dodder (<i>Cuscuta</i> spp.)	Cranberry	Smolder
<i>Bipolaris sorghicola</i>	<i>Sorghum halepense</i>	Cropping systems	-

4. Herbicide-resistant crops: opportunities and concerns

Herbicide-resistant crops allow selective weed control but increase dependency on herbicides. Risks include resistance development, environmental contamination and gene flow to wild relatives. Hence, cautious and integrated use is essential.

5. Integrated Weed Management (IWM)

Integrated Weed Management combines multiple eco-friendly practices to achieve effective and long-term weed control.

- Prevents weed resistance
- Reduces chemical dependency
- Improves crop productivity
- Enhances environmental safety

Traditional practices such as crop rotation, manual weeding and water management, when combined with modern tools, form the backbone of IWM.

Future prospects of eco-friendly weed management

With increasing labour scarcity and climate variability, eco-friendly weed management will become more important.

- Precision tools and mechanization will improve efficiency.
- Greater use of biological agents and natural products is expected.
- Farmer awareness and training will play a key role in adoption.

Sustainable weed management is not just an option but a necessity for future agriculture.

Conclusion

Eco-friendly weed management focuses on working with nature rather than against it. By strengthening crops, improving soil health and promoting biodiversity, weeds can be managed effectively without harming the environment. Adoption of these practices ensures sustainable crop production, safer food and a healthier ecosystem for future generations.

Reference

1. Bahadur, S., Verma, S.K., Prasad, S.K., Madane, A.J., Maurya, S.P., Verma, V.K. and Sihag, S.K. (2015). Eco-friendly weed management for sustainable crop production – A review. *Journal of Crop and Weed*, 11(1): 181-189.
2. Ameta, S.K. and Ameta, S.C. (2021). Eco-friendly approaches of using weeds for sustainable plant growth and production. In: *Plant Performance Under Environmental Stress: Hormones, Biostimulants and Sustainable Plant Growth Management*. Springer International Publishing, Cham, pp. 559-592.

3. Hasan, M., Sarker, M.R.K., Motmainna, M. and Ahmad-Hamdani, M.S. (2024). Bioherbicides: An eco-friendly weed management strategy towards sustainable agriculture. In: *Advances in Tropical Crop Protection*. Springer Nature Switzerland, Cham, pp. 285-304.