

PRODUCTION TECHNOLOGY OF BORONIA (*Boronia Megastigma*): A PREMIUM AROMATIC FLOWER CROP

Rakshitha Y R¹, Chandrashekar S Y² and Bhajantri Archana³

¹PG Scholar, Department of Floriculture and Landscaping, College of Horticulture, Mudigere, India

²Professor and Head, Department of Floriculture and Landscaping, College of Agricultural Sciences, Iruvakk, Shivamogga, India

³Assistant professor, Department of Horticulture, College of Agricultural Sciences, Iruvakk, India

*Corresponding Author Mail ID: rrakshayr@gmail.com

Abstract

Boronia is a high value aromatic flower crop cultivated mainly for its essential oil used in luxury perfumery. The crop grows well in temperate climatic conditions with well-drained soils rich in organic matter. Proper agronomic practices such as selection of quality planting material, nutrient management, irrigation, and careful harvesting are essential for obtaining high-quality flowers and oil. Due to its limited cultivation and high demand, boronia offers excellent scope for income generation in niche markets.

Key words: Boronia, aromatic flower, essential oil, perfumery and high value crop

Introduction

Boronia (*Boronia megastigma*) belongs to the family Rutaceae and is known for its rich, fruity floral fragrance. It is mainly cultivated in regions like Australia and it is highly valued in the perfume industry. The essential oil extracted from its flowers is one of the most expensive natural fragrance ingredients used in luxury perfumes. Due to its high market value and export demand, boronia is gaining importance as a niche aromatic crop.

Botanical description

Boronia is a small evergreen shrub growing up to 1–2 m in height. The plant has slender branches and narrow leaves. The flowers are small, star-shaped and dark purple to brown in color. They possess a strong, complex fragrance that is highly desirable in perfumery.



Types/Varieties

1. ***Boronia megastigma***
 - Most commercially important species
 - High quality essential oil
 - Used in luxury perfumery
2. ***Boronia heterophylla***
 - Ornamental species with pink flowers
 - Limited oil production
3. ***Boronia crenulata***
 - Hardy species
 - Mainly grown for ornamental purposes

Climate and soil requirements

The crop prefers cool temperate climates with temperatures ranging between 10–25°C. It requires moderate rainfall of about 800–1200 mm annually. Well-drained sandy or loamy soils rich in organic matter are ideal. Waterlogging should be avoided as it may lead to root rot.

Propagation and planting material

Propagation is mainly done through semi hardwood cuttings. About 15,000–20,000 cuttings/ha are required for planting.

Healthy, disease free planting materials ensures better establishment and uniform growth. Rooting hormones may be used to improve success rate.

PROPAGATION (CUTTINGS)




Semi-hardwood cuttings of 7–10 cm are taken from healthy mother plants.

Land preparation and planting

The land is prepared thoroughly to ensure good drainage. Raised beds are preferred to avoid water stagnation. Planting is done at a spacing of 1.5 m × 1.5 m, accommodating about 4,400 plants/ha. Proper spacing ensures good aeration and plant development.

LAND PREPARATION



Well drained soil is prepared with raised beds to avoid water stagnation.

PLANTING



Rooted cuttings are planted at a spacing of 1.5 m × 1.5 m.

ROOTING OF CUTTINGS



Cuttings are treated with rooting hormone and placed in rooting media. Roots develop in 4–6 weeks.

VEGETATIVE GROWTH STAGE



Plants grow well under cool climate with regular irrigation and nutrient management.

FLOWERING STAGE



Flowering begins after 1.5–2 years. Flowers are dark purple, small and highly fragrant.

Nutrient management

Application of farmyard manure at 10–15 t/ha during land preparation improves soil fertility. A balanced fertilizer dose of 60–80 kg N, 40–60 kg P₂O₅ and 40–60 kg K₂O per hectare is recommended. Fertilizers are applied in split doses for better nutrient utilization.

Irrigation Management

Light irrigation is provided immediately after planting. Irrigation is given at 7–10 day intervals depending on soil moisture and climatic conditions.

The crop requires moderate moisture levels and over-irrigation should be avoided. Drip irrigation can be adopted for efficient water use.

Weed management

Regular weeding is necessary during early growth stages to reduce competition. Mulching helps conserve moisture and suppress weed growth. Manual and mechanical methods are commonly used. Proper weed control improves plant growth and yield.

Pest and disease management

Boronia is susceptible to root rot and fungal diseases under poor drainage conditions. Common pests include mites and sap sucking insects. Integrated pest management practices such as field sanitation, proper spacing and use of organic sprays help in controlling pests and diseases. Avoiding excess moisture reduces disease incidence.



Harvesting

Flowering generally begins after 1.5–2 years of planting. Flowers are harvested at full bloom stage when fragrance is maximum. Harvesting is done manually due to the delicate nature of flowers. Frequent picking improves overall yield.

Post - harvest processing

Flowers are processed immediately after harvest for essential oil extraction. Solvent extraction is commonly used instead of steam distillation. Proper handling and quick processing help retain fragrance and oil quality.

OIL EXTRACTION (SOLVENT EXTRACTION)



Yield

A well-maintained crop yields about 1.5–2.5 tonnes of flowers per hectare. Essential oil recovery is low, around 0.02–0.05 %, but the oil is highly valuable. Yield depends on plant age, management and environmental conditions.

Conclusion

Boronia is a premium aromatic flower crop with high demand in luxury perfumery. Although cultivation is limited due to specific climatic requirements, it offers excellent economic returns. Adoption of proper cultivation practices and efficient management can enhance productivity and profitability.

References:

1. BROPHY, J. J., GOLDSACK, R. J. AND FORSTER, P. I., 2004, The essential oils of the genus *Boronia* (Rutaceae). *J. Essent. Oil Res.*, **16**(6): 529–536.
2. DUNLOP, P. J., AND BROPHY, J. J., 2007, Chemical composition of *Boronia megastigma* oil. *Flavour Fragr. J.*, **22**(3): 190–193.
3. FAO. 2019, *Aromatic Plants and Essential Oil Crops*. FAO Publications.
4. ICAR. 2020, *Handbook of Floriculture and Landscaping*. ICAR, New Delhi.
5. MACTAVISH, H. S., AND MENARY, R. C., 1997, *Boronia (Boronia megastigma) production and research in Australia*. *Acta Hortic.*, **426**(1): 167–173.
6. PLUMMER, J. A. AND SEDGLEY, M., 1993, Biology and propagation of *Boronia megastigma*. *Acta Hortic.*, **56**(3): 225–234.