



SEXUAL MATURITY, BREEDING SEASON AND GAMETE DEVELOPMENT IN FISHES APPROACHES

Balavinayaga S¹* and Mahalakshmi S²

¹UG Student, Dr. M.G.R Fisheries College and Research Institute, Ponneri-601 204, Tamil Nadu Dr. J. Jayalalitha Fisheries University, Tamil Nadu, India.

²PG Scholar, Department of Aquatic Environment Management, Dr. M.G.R Fisheries College and Research Institute, Ponneri-601 204, Tamil Nadu Dr. J. Jayalalitha Fisheries University, Tamil Nadu, India.

*Corresponding Author Mail ID: balavinayaga023@gmail.com

Abstract

Sexual maturity in fishes is a critical biological process that determines their ability to reproduce and sustain populations. It involves the development of reproductive organs and the production of functional gametes, namely eggs in females and sperm in males. This process, known as gonadal maturation, is influenced by factors such as age, size, environmental conditions, and hormonal regulation. Environmental parameters like temperature, rainfall, and photoperiod play a significant role in initiating and regulating maturation and breeding cycles. Fishes exhibit distinct breeding patterns, with many freshwater species, particularly major carps, breeding during the south-west monsoon, while some species like tilapia and common carp are capable of breeding throughout the year. Marine and brackish water fishes often have extended breeding seasons. Gamete development occurs through well-defined stages, including oogenesis in females and spermatogenesis in males, both of which are regulated by hormones such as gonadotropins, Oestrogen, and testosterone. Hormonal control is essential for coordinating the growth of gonads, maturation of gametes, and spawning processes. The interaction between environmental cues and endocrine mechanisms ensures successful reproduction under favourable conditions. Understanding these processes is highly important in aquaculture, as it aids in effective broodstock management, induced breeding, and improved seed production. Overall, knowledge of

sexual maturity and reproductive biology in fishes contributes to sustainable fishery practices and enhanced aquaculture productivity.

Introduction

Sexual maturity in fishes refers to the stage in the life cycle when a fish becomes capable of reproduction. At this stage, the reproductive organs, namely the testes in males and ovaries in females, become fully developed and are able to produce viable gametes such as sperm and eggs. The process leading to sexual maturity involves the gradual development of immature germ cells into functional reproductive cells through a series of biological and hormonal changes. This stage is very important because only sexually mature fishes can participate in spawning and contribute to the continuation of the species. Sexual maturity plays a crucial role in aquaculture and fishery management. In aquaculture, the successful breeding of fishes depends largely on the proper identification and management of mature broodstock. Farmers need to know when fishes attain maturity so that they can plan induced breeding, seed production, and stocking efficiently. Understanding sexual maturity helps in increasing fish production, improving seed quality, and ensuring a continuous supply of fish for commercial purposes. It also helps in conserving natural fish populations by preventing overfishing of immature individuals and allowing them to reproduce before being harvested. The age at which fishes attain sexual maturity varies widely among different species.

Some fishes, such as tilapia, reach maturity within a few months, while others, like major carps, may take one to two years or more. Size is another important factor, as fishes generally need to attain a certain body length or weight before becoming mature. In many cases, even if a fish has reached the appropriate age, it may not become sexually mature unless it has achieved the required size. Environmental factors also have a significant influence on sexual maturity in fishes. Parameters such as water temperature, rainfall, photoperiod (day length), and availability of food play a vital role in regulating the maturation process. For instance, many freshwater fishes in tropical regions attain maturity and breed during the monsoon season when environmental conditions are favourable. Changes in these environmental conditions can either accelerate or delay the onset of maturity. In addition to these factors, hormonal regulation within the fish body controls the maturation process. Hormones coordinate the development of gonads and the production of gametes, ensuring that fishes become reproductively active at the right time. Thus, sexual maturity in fishes is a complex process influenced by biological, environmental, and hormonal factors, and it forms the foundation for successful reproduction and sustainable aquaculture practices.

Sexual Maturity

Sexual maturity in fishes is closely associated with the process of gonadal maturation, which refers to the development and functional readiness of the reproductive organs testes in males and ovaries in females. Gonadal maturation is a complex biological process in which immature gonads gradually develop into fully functional organs capable of producing and releasing gametes. During this process, small and undeveloped germ cells grow, multiply, and differentiate into mature reproductive cells, ultimately enabling the fish to participate in reproduction. The development of the gonads involves significant structural and physiological changes. In female fishes, the ovaries increase in size and weight as the oocytes (egg cells) develop and accumulate yolk, which serves as a source of nutrition for the developing embryo.

The ovary may occupy a considerable portion of the body cavity when fully mature. In male fishes, the testes also undergo enlargement and become filled with spermatozoa. The testes are typically creamy white in appearance when mature and are capable of producing large quantities of sperm. The production of gametes is a key outcome of gonadal maturation. In females, this process is known as oogenesis, where oogonia develop into mature ova (eggs) through several stages of growth and maturation. In males, the process is called spermatogenesis, where spermatogonia develop into sperm cells. These processes are regulated by hormones, especially gonadotropins, which control the growth, development, and final maturation of the gonads. Once the gametes are fully developed, they are released during spawning for fertilization. There are several important characteristics of sexual maturity in fishes. One notable feature is that males generally attain sexual maturity earlier than females. This difference may be due to variations in energy requirements and reproductive roles between the sexes. Additionally, the age and size at which fishes reach maturity vary greatly among species. Some species mature rapidly within a few months, while others may take several years to become reproductively active.

Environmental factors play a significant role in influencing gonadal maturation and sexual maturity. Conditions such as water temperature, photoperiod (length of daylight), rainfall, and availability of food can either accelerate or delay the maturation process. Favourable environmental conditions often trigger hormonal changes that promote the development of gonads and the production of gametes. Conversely, unfavourable conditions may suppress or postpone maturation. Gonadal maturation is a vital process that leads to sexual maturity in fishes, involving the development of reproductive organs and the production of gametes.

It is influenced by a combination of biological, environmental, and hormonal factors, and it plays a key role in the reproductive success and sustainability of fish populations.



Breeding Season in Fishes


The breeding season in fishes refers to the specific period during which fishes become reproductively active and spawn. Unlike many terrestrial animals, fishes often show strong seasonal patterns in their breeding behaviour, which are closely linked to environmental conditions. This seasonal breeding ensures that the survival of eggs, larvae, and juveniles is maximized by synchronizing reproduction with favourable environmental conditions such as food availability and suitable habitat. In many freshwater fishes, especially in tropical countries like India, breeding is closely associated with the south-west monsoon season. Major carps such as Catla, Rohu, and Mrigal typically attain maturity and breed during this period. The monsoon brings a sudden increase in water level, flow, and nutrient availability in rivers, ponds, and floodplains, creating ideal conditions for spawning and the growth of offspring. These environmental changes act as natural stimuli that trigger reproductive activities in fishes. However, not all fishes follow strict seasonal breeding patterns. Some species, such as tilapia and common carp, are capable of breeding throughout the year under favourable conditions. These fishes are known as continuous or multiple spawners, as they can reproduce several times within a year. Their ability to breed continuously makes them highly suitable for aquaculture, as it ensures a constant supply of fish seed. Marine and brackish water fishes often exhibit comparatively longer and more flexible breeding seasons. Many of these species begin their reproductive activities during the pre-monsoon period and continue through the monsoon and even beyond. The extended breeding period in

such fishes increases the chances of successful fertilization and survival of larvae in dynamic aquatic environments. Environmental factors play a crucial role in regulating the breeding season in fishes. Temperature is one of the most important factors, as it influences metabolic activities, hormone production, and gonadal development. Optimal temperature ranges are necessary for the initiation of spawning. Rainfall is another key factor, particularly in freshwater ecosystems, where it leads to increased water levels, reduced salinity, and improved nutrient availability, all of which favour breeding. Additionally, photoperiod (day length) acts as an important environmental cue that helps synchronize reproductive cycles. Changes in day length are detected by fishes and influence hormonal mechanisms that control maturation and spawning. The breeding season in fishes is largely determined by environmental conditions and varies among species. While some fishes breed seasonally, others reproduce throughout the year. Understanding these patterns is essential for effective fishery management and aquaculture practices, as it helps in planning breeding programs and ensuring successful seed production.

Gamete Development in Females (Oogenesis)

Oogenesis is the process by which female gametes (eggs or ova) are formed and developed in the ovaries of fishes. It is a complex and continuous process involving the transformation of primitive germ cells into fully mature eggs capable of fertilization. This process occurs in several well-defined stages, each characterized by specific structural and physiological changes. Oogenesis is regulated by both environmental factors and hormones, ensuring proper timing of maturation and spawning.

Gamete Development in Female Fishes



A. Oogenesis (Formation of Oogonia to Oocytes)

The process of oogenesis begins with the formation of oogonia, which are the primary germ cells present in the ovary. These cells divide and differentiate to form oocytes through a process involving meiosis (reduction division). Oogenesis occurs throughout the life of many fishes, especially in species that spawn multiple times. The continuous presence of developing oocytes ensures that fishes can reproduce repeatedly under suitable conditions.

B. Primary Growth Stage

During the primary growth stage, the oocytes increase significantly in size due to the accumulation of cellular components such as cytoplasm and organelles. At this stage, the oocyte becomes surrounded by a layer of follicular cells, forming a structure known as the follicle. Additional layers, such as the zona pellucida (a protective layer) and theca, also develop around the oocyte. This stage is generally independent of hormonal control and represents the initial phase of oocyte growth. By the end of this stage, the oocyte is referred to as a pre-vitellogenic oocyte.

C. Yolk Vesicle Formation

The next stage involves the formation of yolk vesicles within the cytoplasm of the oocyte. This stage is triggered by environmental factors such as changes in temperature, rainfall, and photoperiod. Hormonal stimulation, particularly by gonadotropins (GtH), also plays a role in initiating this process. The yolk vesicles contain glycoproteins and serve as precursors to energy reserves that will support the embryo after fertilization. This stage is sometimes referred to as early or endogenous vitellogenesis.

D. Vitellogenesis

Vitellogenesis is the most important and major growth phase of oocyte development. During this stage, large amounts of yolk proteins (vitellogenin) are synthesized in the liver under the influence of the hormone Oestrogen. These proteins are transported through the bloodstream to the ovary, where they are taken up by the oocytes and stored as yolk. This leads to a rapid


increase in the size and weight of the oocyte. Hormones such as gonadotropins (GtH) and Oestrogen play a crucial role in regulating this stage. Vitellogenesis ensures that the developing egg contains sufficient nutrients for the growth of the embryo.

E. Final Maturation

Final maturation is a rapid stage that prepares the oocyte for ovulation. It involves the migration of the germinal vesicle (nucleus) towards the periphery of the oocyte, followed by its breakdown (GVBD – germinal vesicle breakdown). This process marks the resumption and completion of meiosis. Final maturation usually occurs within a short period, often less than 24 hours in many fish species. During this stage, hormonal changes occur, including a shift from Oestrogen production to maturation-inducing steroids, which trigger the final steps of egg development.

F. Ovulation


Ovulation is the final stage of oogenesis, during which the fully mature egg is released from the follicle into the ovarian cavity. The egg is then ready to be spawned into the external environment for fertilization. This process is controlled by hormones, particularly prostaglandins, which are produced under the influence of gonadotropins. After ovulation, the eggs remain viable for a limited period, after which they may become overripe and lose their ability to be fertilized.



Gamete Development in Males

The development of male gametes in fishes is known as spermatogenesis, which occurs in the testes. This process involves the transformation of primitive germ cells into mature spermatozoa capable of fertilizing eggs. Compared to females, male fishes generally attain maturity earlier and remain reproductively active for a longer period. The process of gamete development in males is continuous and consists of several stages, each regulated by hormones and influenced by environmental factors.

Gamete Development in Male Fishes



A. Spermatogenesis

Spermatogenesis is the initial stage in the development of male gametes. It begins with spermatogonia, which are undifferentiated germ cells present in the testes. These cells undergo repeated divisions and eventually differentiate into spermatocytes through the process of meiosis. This stage is primarily controlled by the hormone testosterone, which is produced by specialized cells known as Leydig cells under the influence of gonadotropins (GtH). During this stage, the developing cells are supported and nourished by Sertoli cells within the testes.

B. Spermiogenesis

Spermiogenesis is the next stage, during which the spermatocytes further develop into spermatids and then transform into mature sperm cells. This stage involves significant structural changes, such as the formation of the head, middle piece, and tail (flagellum) of the sperm. The development of the flagellum enables the sperm to become motile. Unlike earlier stages, spermiogenesis can occur even in the absence of

strong hormonal influence and mainly focuses on the maturation and differentiation of sperm cells.

C. Spermiation

Spermiation is the process by which fully developed spermatozoa are released from the seminiferous tubules into the lumen of the testes. This stage is regulated by hormones such as 11-ketotestosterone, which is produced by Leydig cells. During spermiation, sperm cells are freed from the supporting Sertoli cells and become available for release during spawning. This process ensures that a sufficient quantity of mature sperm is ready for fertilization at the appropriate time.

D. Hydration

Hydration is the final step in the development and release of male gametes. In this stage, spermatozoa are mixed with a fluid secreted by the walls of the sperm duct, resulting in the formation of milt, which is the fluid containing sperm. This dilution process is essential for the effective release and movement of sperm during spawning. Hydration is influenced by hormones such as 17α -20 β -dihydroxyprogesterone and is often synchronized with the final maturation and ovulation in females to ensure successful fertilization.

Hormonal Control

Hormonal control plays a vital role in regulating reproduction in fishes. The processes of gonadal growth, maturation, and spawning are not random but are carefully coordinated by a complex system of hormones. These hormones are produced by different organs such as the brain, pituitary gland, and gonads, and they act together to ensure proper timing and success of reproduction. Environmental factors like temperature, rainfall, and photoperiod influence this hormonal system, triggering the release of specific hormones at the right time.

Role of Gonadotropins (GtH)

Gonadotropins (GtH) are the primary hormones responsible for controlling reproductive processes in fishes. They are secreted by the pituitary gland and play a central role in regulating the development and functioning of gonads.

GtH stimulates the testes and ovaries to grow and produce sex hormones. In females, it promotes the development of oocytes and supports processes such as vitellogenesis and final maturation. In males, GtH stimulates the production of testosterone and other androgens, which are essential for spermatogenesis. Thus, GtH acts as a key regulator that initiates and coordinates all stages of reproduction.

Role of Oestrogen

Oestrogen is a major female sex hormone produced mainly in the ovaries. It plays a crucial role in the development and maturation of oocytes. One of its primary functions is to stimulate the liver to produce vitellogenin, a yolk protein precursor. This vitellogenin is transported through the bloodstream to the oocytes, where it is deposited as yolk, providing essential nutrients for the developing embryo. Oestrogen also helps in the growth of ovarian tissues and regulates feedback mechanisms that control the release of gonadotropins.

Role of Testosterone

Testosterone is the primary male sex hormone produced in the testes by Leydig cells. It is essential for the development of male reproductive organs and the process of spermatogenesis. Testosterone promotes the division of spermatogonia into spermatocytes and supports the overall growth and maturation of testes. It also influences secondary sexual characteristics and reproductive behaviour in fishes. In addition, testosterone may be converted into other hormones that play roles in final maturation and spawning.

Hormonal Regulation of Reproductive Processes

Growth of Gonads: Hormones like gonadotropins stimulate the enlargement and development of ovaries and testes, preparing them for reproductive activity.

Maturation: Hormonal interactions control the development of gametes, including oocyte growth in females and sperm formation in males. Oestrogen and testosterone play important roles in this stage.

Spawning: The final release of eggs and sperm is also hormonally controlled. Gonadotropins trigger the production of other hormones such as maturation-inducing steroids and prostaglandins, which lead to ovulation in females and spermiation in males.

Conclusion

Sexual maturity is a fundamental aspect of fish reproduction, as it determines the ability of fishes to produce viable gametes and successfully participate in spawning. The development of reproductive organs and the formation of eggs and sperm are essential processes that ensure the continuation of fish species. Without proper attainment of sexual maturity, reproduction cannot occur, making it a critical stage in the life cycle of fishes. The process of sexual maturity is influenced by a combination of environmental and hormonal factors. Environmental conditions such as temperature, rainfall, and photoperiod play a significant role in triggering maturation and breeding activities.

At the same time, hormones like gonadotropins, estrogen, and testosterone regulate the growth of gonads, development of gametes, and the timing of spawning. The interaction between these internal and external factors ensures that reproduction occurs under favorable conditions for the survival of offspring. Understanding sexual maturity and its controlling factors is highly important in aquaculture practices. It helps fish farmers manage broodstock effectively, plan breeding programs, and improve seed production. Knowledge of maturation and spawning cycles also supports the use of techniques such as induced breeding, leading to increased fish production and sustainability.

Therefore, proper understanding of sexual maturity not only enhances reproductive success but also contributes significantly to the development of efficient and profitable aquaculture systems.

References

1. Wootton, R. J., & Smith, C. (2015). Reproductive biology of teleost fishes. Wiley-Blackwell.
2. Jakobsen, T., Fogarty, M. J., Megrey, B. A., & Moksness, E. (2009). Fish reproductive biology: Implications for assessment and management. Wiley-Blackwell.
3. Yoshida, M., & Asturiano, J. F. (2020). Reproduction in aquatic animals: From basic biology to aquaculture technology. Springer.
4. Sundaray, J. K., Rather, M. A., Kumar, S., & Agarwal, D. (2021). Molecular endocrinology and reproductive physiology of fish. Springer.
5. Lee, C. S., & Donaldson, E. M. (2001). Reproductive biotechnology in finfish aquaculture. Elsevier.
6. Chen, X., & Liu, B. (2022). Biology of fishery resources. Springer.
7. Zohar, Y. (2020). Fish reproductive biology and endocrinology. General and Comparative Endocrinology.